

A New Detached M Dwarf Eclipsing Binary

O.L. Creevey^{1,2,3}, G.F. Benedict⁴, T.M. Brown¹, R. Alonso³, P. Cargile⁴, G. Mandushev⁵,
D. Charbonneau⁶, B.E. McArthur⁴, W. Cochran⁴, F.T. O'Donovan⁷, S.J. Jiménez-Reyes³,
J.A. Belmonte³, D. Kolinski¹

creevey@hao.ucar.edu

fritz@astro.as.utexas.edu

timbrown@hao.ucar.edu

ras@iac.es

p.cargile@mail.utexas.edu

gmand@lowell.edu

dcharbonneau@cfa.harvard.edu

mca@barney.as.utexas.edu

wdc@shiraz.as.utexas.edu

francis@caltech.edu

sjimenez@iac.es

jba@iac.es

kolinski@hao.ucar.edu

ABSTRACT

¹High Altitude Observatory, National Center for Atmospheric Research, Boulder 80307, CO USA

²Universidad de La Laguna, 38206 La Laguna, Tenerife, Spain

³Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain

⁴McDonald Observatory, University of Texas, Austin, TX 78712 USA

⁵Lowell Observatory, Flagstaff, AZ 86001 USA

⁶Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138 USA

⁷California Institute of Technology, 1200 E. California Blvd., Pasadena, CA 91125 USA

We describe a newly-discovered detached M-dwarf eclipsing binary system. This system was first observed by the TrES network during a long term photometry campaign of 54 nights. Analysis of the folded light curve indicates two very similar components orbiting each other with a period of 1.12079 ± 0.00001 days. Spectroscopic observations with the Hobby-Eberly Telescope show the system to consist of two M3e dwarfs in a near-circular orbit. Double-line radial velocity amplitudes, combined with the orbital inclination derived from light-curve fitting, yield $M_{total} = 0.983 \pm 0.007 M_{\odot}$, with component masses of $M_1 = 0.493 \pm 0.003 M_{\odot}$ and $M_2 = 0.489 \pm 0.003 M_{\odot}$. The light-curve fit yields component radii of $R_1 = 0.453 \pm 0.060 R_{\odot}$ and $R_2 = 0.452 \pm 0.050 R_{\odot}$. Though a precise parallax is lacking, broadband VJHK colors and spectral typing suggest component absolute magnitudes of $M_V(1) = 11.18 \pm 0.30$ and $M_V(2) = 11.28 \pm 0.30$.

Subject headings: binaries: eclipsing—binaries: close—stars: late-type—stars: individual: (TrES-Her0-07621)

1. Introduction

Although low mass binary stars are the most abundant stars in the galaxy (Henry et al. 1999), their intrinsic faintness inhibits their detection and study. Non-contact eclipsing binary M dwarf systems have great value, as these systems allow accurate estimates of the most basic stellar parameters: mass and radius. Only four¹ such systems are known and have been studied in detail; YY Gem (Bopp 1974; Leung & Schneider 1978), CM Dra (Lacy 1977; Metcalfe et al. 1996; Kozhevnikova et al. 2004), GJ 2069A (Delfosse et al. 1999; Ribas 2003), and OGLE BW03 V038² (Maceroni & Montalbán, 2004). The observed properties of each of these systems present discrepancies with the theory of low-mass stellar objects; neither the observed mass-radius nor mass-luminosity relations are well represented by existing models (Benedict 2000); see Figure 1. The problem most likely lies in the shortcomings of the physical models, owing to the lack of understanding of the complex atmospheres of such low-mass objects (Baraffe et al. 1998). Enlarging the small existing sample of such systems is therefore desirable, to allow more detailed comparisons between observations and the theory of these ubiquitous, interesting, and complex objects. Here we report preliminary analysis of a fourth such low-mass eclipsing binary.

¹We refer specifically to binaries where both components are M Dwarfs. There have been, however, a number of M stars whose companion is an F or G MS star (e.g. Pont et al., 2004), see Figure 1

²This is a very close although still detached system.

2. Observations

2.1. Photometric Observations

The recently discovered spectroscopic binary, TrES–Her0-07621 ($\alpha=16^h50^m20.7^s$, $\delta=+46^\circ39'01''$ (J2000), $V=15.51 \pm 0.08$) was first identified through an analysis of photometric time series from the TrES (Trans-Atlantic Exoplanet Survey) network. This network consists of three telescopes: *STellar Astrophysics and Research on Exoplanets* (STARE,³ Brown & Charbonneau, 1999), *Planet Search Survey Telescope* (PSST⁴, Dunham et al. 2004), and *Sleuth*⁵. The telescopes are similar in their characteristics, with apertures of 10cm, 2048×2048 pixel CCD detectors and fields of view of $6^\circ \times 6^\circ$.

TrES collects long-term time-series photometry in one filter. The photometry run in question spanned 54 days, beginning May 6 2003, and was observed in a band roughly equivalent to Harris R at a cadence of 1 image every 2 minutes. The images were reduced and calibrated by an automatic package developed specifically for these data. TrES–Her0-07621 was observed by both STARE and PSST, but the latter time series proved significantly noisier. We therefore analyzed only the STARE lightcurve. The R magnitude is 14.42 with each point having a formal accuracy of 0.04 mag rms. This lightcurve contains 8781 data points, obtained in 309.5 hours over 54 days, giving a duty cycle of 23.8%⁶.

A high-SNR peak in the time series’ frequency spectrum at 1.79 cycles per day initiated the study of TrES–Her0-07621. Folding the star’s light curve with a period of 1.1208 d showed it to be an eclipsing binary. The light curve also displays sinusoidal out-of-eclipse variations near the photometric period. The star’s infrared colors from the 2MASS⁷ catalog are quite red, (Table 1), and the USNO-B⁸ catalog shows a significant proper motion. Taken together, these facts suggested that the object is a binary M dwarf, with substantial levels of magnetic activity driven by the rapid, tidally-locked rotation of the component stars; this motivated further study.

³Observatorio del Teide, Tenerife, Spain

⁴Lowell Observatory, AZ, USA

⁵Palomar Observatory, CA, USA. <http://www.astro.caltech.edu/~ftod/tres/sleuth.html>

⁶The data are available via STARE website <http://www.hao.ucar.edu/public/research/stare/stare.html>

⁷Two Micron All Sky Survey: University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology <http://irsa.ipac.caltech.edu/cgi-bin/Gator/nph-dd>

⁸SIMBAD, operated at CDS, Strasbourg, France; the NASA/IPAC Extragalactic Database (NED) and supported by JPL, California Institute of Technology, <http://www.nofs.navy.mil/data/fchpix/>

2.2. Spectroscopic Observations

In September 2004 we obtained spectroscopic observations of TrES–Her0-07621 using the High Resolution Spectrograph (HRS, Tull 1998) on the Hobby-Eberly Telescope (HET). We secured measurements at 4 epochs; each epoch contained three separate exposures taken over approximately one hour — giving a total of 12 spectra. The analysis was carried out with standard IRAF (Tody 1993) `echelle` and `rv` package tools, including `fxcorr`. We cross-correlate TrES–Her0-07621 with an M2 dwarf (Gl 623) template and extract velocities for both components at four distinct phases. We adopted a radial velocity for the Gl 623 primary of -29.2 km s^{-1} , given the orbital phase at which the template was secured and a systematic velocity, $V_{sys} = -27.5 \text{ km s}^{-1}$, from Marcy & Moore (1989). The HRS utilizes two CCDs covering the blue and red spectral regions. The data from each chip were analyzed independently, resulting in two velocity estimates. A third velocity estimate was obtained by cross-correlating an artificial H-alpha emission template with the H-alpha emission line found in each exposure. Given the large orbital velocities, there was no blending of correlation peaks at any phase. The three velocities (blue, red, H α) are obviously not independent determinations, but do provide an estimate of our internal error.

3. Analysis

Figure 2 shows the component velocities plotted against photometric phase, while figure 3 (top panel) shows the folded photometric light curve. It is evident from the nearly-symmetrical and sinusoidal radial velocity variation and from the highly symmetrical light curve that the orbit is nearly circular, and that the component masses and surface brightnesses are similar.

An initial period analysis of the entire STARE lightcurve using the technique of phase dispersion minimization refined the photometric period to 1.1209 ± 0.0006 days. We predicted and then observed an eclipse on 14 May 2004 using the 1.2m telescope at the Fred L. Whipple Observatory, AZ, USA using SLOAN filters r , i and z . The long time base provided by this observation allowed us to refine the photometric period. By fitting the lightcurves during (21) eclipse times (we included only totally observed eclipses) to parabolas, we determined all the times of minimum light (center of eclipse) with corresponding error. For the eclipse observed on May 14th 2004, we only used the time of minimum light from the r filter. We also used observations from IAC80 (see below). Using the bootstrap method we refined the period to 1.12079 ± 0.00001 corresponding to a precision of 1 second. The epoch of secondary minimum, T_0 , was meanwhile determined to be 2453139.749509 (HJD) ± 0.000075 .

TrES–Her0-07621 has a stellar neighbor at a distance of 8", close enough that the two objects are blended in our STARE observations (STARE has a pixel size of about 11 arcsec). Observations in *R* and *I* Johnson filters using the IAC80 at Observatorio del Teide on 30 August 2004 provided a more realistic picture of the depth of one of the eclipses, while also allowing us to confirm the photometric period. We measured the PSF of both the binary and the neighbor using all five images outside of the eclipse. From these we derived the *R* fractional flux contribution from this companion star of 0.19 ± 0.04 . Because the companion star is also quite red (Table 1), the flux should be similar (to within the error) in both Johnson and Harris *R* filters, and so we can use this number to analyze the STARE time series. Measurement of the contamination of the eclipse signal from TrES–Her0-07621 by the companion star is important, because it must be accounted for when fitting the time-series data to estimate the stellar radii. This neighbor also has a proper motion that is similar in magnitude and direction to that of TrES–Her0-07621, indicating the possibility that TrES–Her0-07621 is at least a triple system, with the eclipsing pair of stars accompanied by a third M dwarf at a distance of hundreds of AU.

Adopting the photometric period as the orbital period and introducing its associated error, we fit all 36 radial velocities (blue, red, H α) with a Keplerian model using GaussFit (Jefferys, Fitzpatrick & McArthur, 1988). The model is similar to that used in McArthur et al. (2004). We assume an eccentricity e of 0. The resulting radial velocity semi-amplitudes are $K_1 = 100.54 \pm 0.31 \text{ km s}^{-1}$ and $K_2 = 101.29 \pm 0.31 \text{ km s}^{-1}$, $(M_1 + M_2) \sin^3 i = 0.9547 \pm 0.0062 M_\odot$, and $M_1/M_2 = 1.0075 \pm 0.0044$. A formal solution including eccentricity ($e = 0.006 \pm 0.002$) provided a better solution, reducing χ^2 by 8%, while reducing the number of degrees of freedom by 3%. However, we constrain $e = 0$ for this analysis.

We developed a chi-square minimization algorithm to estimate orbital parameters from the light curve, ignoring any variations between eclipses (Figure 3). The input parameters are period P , component masses M_1 and M_2 , limb-darkening coefficients (0.7, Claret, 1998, Table 7) and the light from a third nearby star as a fraction of the total light of the system (0.19 ± 0.04). The code solves for both radii R_1 and R_2 , effective temperature ratio T_2/T_1 , center of minimum eclipse T_0 and inclination i . Figure 3 shows the resulting fit to the data.

The initial estimates for R_1 , R_2 , T_2/T_1 and i were derived from two-dimensional χ^2 contour plots (while keeping the other two parameters fixed). These contour plots presented high correlations between the two radii, constraining their sum while insensitive to their difference; and between radius (R_1 or R_2) and inclination; larger radius implies smaller inclination. T_2/T_1 was uncorrelated to both radii and inclination, so its error is given by the corresponding value of this parameter at $\chi^2 + \sigma$ (Press et al, 1986) in the direction of its axis. However, because the other parameters are obviously not independent, R_1 and R_2

for example, the error spanned the range of radii where the contour value is $\chi^2 + \sigma$, (the full range error ellipsoid).

Even with the component masses determined, in absence of a T_{eff} measurement we require the component absolute magnitudes to place these stars on the Mass-Luminosity Relation (MLR). From the TrES data, calibrated using stars within 1° that have measured V magnitudes from SIMBAD, we obtain a V-band apparent magnitude of 15.51 ± 0.08 for the combined 3-star system, giving $V-K = 4.63 \pm 0.10$ (Table 1). Assuming a wavelength independent relative flux we estimate $\Delta V_{AB-C} = 1.72$ from difference J, H and K magnitudes (between the neighbor and the binary). We can also estimate ΔV_{A-B} (between the binary components) $= 0.1 \pm 0.05$, based on the derived temperature and radii differences. Taking all of the above into account, we estimate the component magnitudes of $V_A = 16.37 \pm 0.1$, $V_B = 16.56 \pm 0.1$ and $V_C = 17.43 \pm 0.1$, where C is the stellar neighbor.

From Hawley et al, (2002) color-spectral type relations we estimate an M3 spectral type for each component. We obtain from the Hawley M_J - spectral type relationships component absolute magnitudes of $M_V = 11.18, 11.28 \pm 0.3$. Accepting this estimate of the luminosities, the distance modulus is μ ($\sim 16.4-11.2$) ~ 5.2 , corresponding to $d \sim 110$ pc. For this nearby system we have assumed no absorption ($A_V = 0$). We also use the radii and effective temperature (Table 3) to determine luminosities, differentially with respect to the Sun (e.g. Benedict et al, 2003). With bolometric corrections as a function of temperature from Flower (1996) we obtain an average $d = 118 \pm 13$ pc for the two components.

4. Results and Comments

Using all the derived parameters and errors, we refit the lightcurve using our code, and tested these results with the code *Nightfall*⁹ (see below). Both codes give similar results, their difference being within the error bars. Tables 2 and 3 summarize the results.

Our code does not allow for stellar spots, so we subtracted a smooth function (by a Fourier technique) to remove the out-of-eclipse variations making a rectified lightcurve. We also constrain $e = 0$. The top panel of Figure 3 shows the synthetic lightcurve (continuous line) corresponding to the model fit (our code) of the folded light curve (small crosses). Phase = 0 corresponds to the secondary eclipse. The bottom panel shows the residuals of the fit. The residuals show no variation as a function of phase indicating an adequate model fit.

Because our code is unable to account for spot variability, we inspected the residuals

⁹<http://www.lsw.uni-heidelberg.de/users/rwichman/Nightfall.html>

after subtracting the model fit from the unrectified lightcurve. These residuals also showed no evidence of eclipses. We also fit this unrectified lightcurve to find R_1 , R_2 , i , T_0 and T_2/T_1 . The results varied slightly from those for the rectified light curve, but stayed within the error bars (Table 3).

Our original (unrectified) photometric light curve contains non-uniform outside-eclipse variations. Binary systems such as TrES–Her0-07621 are often magnetically active (e.g., Strassmeier et al. 1993). While tidal effects may be important, these non-uniform variations are most likely explained by star spots. We used *Nightfall* to model our unrectified lightcurve, because this code allows for the presence of spots on each of the components. Our derived parameters were used as inputs and we attempted to solve for the longitude, latitude and radii of spot(s). There was no unique solution; many combinations of these spot parameters could compensate for the out-of-eclipse variations, although they always presented a 180° longitude difference. This preferred longitude difference has also been observed in other active binary systems (see e.g., Henry et al. 1995). The presence of spots can have a significant effect on the accuracy of the derived parameters, such as inclination, temperature and radii (Torres & Ribas (2002) discuss this for the case of YY Gem). Additional observations, photometry in particular, will be necessary to increase the precision of the radii estimates as well as to learn more about the magnetic behaviour of the stars. This could then provide a link towards a better understanding of the physical processes of these low-mass objects.

We thank Hectór Vazquez Ramio (observations on IAC80), and operating staff for STARE. The IAC80 and STARE are operated by the Instituto de Astrofísica de Canarias in the Spanish Observatorio del Teide. We thank Mark Everett (observations on 48-inch telescope at Fred L. Whipple Observatory on Mount Hopkins, Arizona USA, operated by Harvard-Smithsonian Center for Astrophysics.) We also thank Rainer Wichmann for the use of the program for the light-curve synthesis, *Nightfall*. Support for this work was provided by NASA through grants GO-09408 and GO-09407 the Space Telescope Science Institute, which is operated by the Association of Universities of Research in Astronomy, Inc., under NASA contract NAS5-26555. The Hobby-Eberly Telescope (HET) is a joint project of the University of Texas at Austin, the Pennsylvania State University, Stanford University, Ludwig-Maximilians-Universität München, and Georg-August-Universität, Göttingen. We thank the HET resident astronomers and telescope operators. We thank the referee for their constructive comments.

REFERENCES

- Baraffe, I., Chabrier, G., Allard, F. & Hauschildt, P.H. 1998, *A&A*, 337, 403
- Benedict, G.F., McArthur, B.E., Franz, O.G., Wasserman, L.H., Henry, T.J. 2000, *AJ*, 120, 1106
- Benedict, G. F., et al. 2003, *AJ*, 126, 2549
- Bopp, B.W. 1974, *ApJ*, 193, 389
- Bouchy, F., Pont, F., Melo, C., Santos, N. C., Mayor, M., Queloz, D., & Udry, S. 2005, *A&A*, 431, 1105
- Brown, T. M. & Charbonneau, D. 1999, *Bulletin of the American Astronomical Society*, 31, 1534
- Claret, A. 1998, *A&A*, 335, 647
- Cochran, W.D., et al. 2004, *ApJ*, L611
- Cox, A.N. 2000, *Allen’s Astrophysical Quantities* (4th ed.; New York, AIP Press: Springer)
- Delfosse, X., Forveille, T., Mayor, M., Burnet, M., & Perrier, C. 1999, *A&A*, 341, L63
- Dunham, E. W., Mandushev, G. I., Taylor, B. W., & Oetiker, B. 2004, *PASP*, 116, 1072
- Flower, P. J. 1996, *ApJ*, 469, 355
- Hawley, S.L., et al. 2002, *AJ*, 123, 3409
- Henry, T. J., & McCarthy, D. W. 1993, *AJ*, 106, 773
- Henry, G.W., Eaton, J.A., Hamer, J., & Hall, D.S. 1995, *ApJS*, 97, 513
- Jefferys, W. H., Fitzpatrick, M. J., & McArthur, B. E. 1988, *Celestial Mechanics*, 41, 39
- Kozhevnikova, A.V., Kozhevnikov, V.P., Zakharova, P.S., Polushina, T.S., Svechnikov, M.A. 2004, *Astronomy Reports*, 48, 826
- Lacy, C. 1977, *ApJ*, 218, 444
- Leung, K., & Schneider, D. 1978, *AJ*, 83, 618
- Maceroni, C., & Montalbán, J. 2004, *A&A*, 426, 577

- Marcy, G.W., & Moore, D., 1989, ApJ, 341, 961
- McArthur, B.E., et al. 2004, ApJ, L614, L81
- Metcalfe, T.S., Mathieu, R.D., Latham, D.W., Torres, G. 1996, ApJ, 456, 356
- Pont, F., Bouchy, F., Melo, C., Santos, N.C., Mayor, M., Queloz, D., Udry, S. 2005, astro-ph/0501615
- Press, W.H., Flannery, B.P., Teukolsky, S.A., Vetterling, W.T., 1986, Numerical Recipes, Cambridge, Sect 14.5
- Ribas, I. 2003, A&A, 398, 239
- Strassmeier, K. G., Hall, D. S., Fekel, F. C., & Scheck, M. 1993, A&AS, 100, 173
- Torres, G., & Ribas, I. 2002, ApJ, 567, 1140
- Tody, D. 1993, ASP Conf. Ser. 52: Astronomical Data Analysis Software and Systems II, 2, 173
- Tull, R.G. 1998, Proc. SPIE, 3355, 387

Table 1: Catalog Information

	Binary	Neighbor
J ^a	11.773	13.487
H	11.137	12.863
K	10.880	12.615
J-K	0.893	0.872
μ_{α} ^b	-2 ± 8	-26 ± 14
μ_{δ}	$+30 \pm 3$	$+28 \pm 7$

^a2MASS catalog

^bUSNO-B catalog, units mas yr⁻¹

Table 2: System Parameters

P(days)	1.12079	$\pm$	0.00001
P(years)	0.00306861	$\pm$	0.00000036
T ₀ (JHD)	2453139.749509	$\pm$	0.00075
M _T (M _☉)	0.983	$\pm$	0.007
a(AU)	0.01047	$\pm$	0.00002
a(R _☉)	2.251	$\pm$	0.005
<i>i</i> (°)	83.12	$\pm$	0.30
γ(km s ⁻¹)	-26.5	$\pm$	0.3
T _B /T _A	0.97	$\pm$	0.02

Table 3: Component Parameters

	A			B		
$M(M_{\odot})$	0.493	$\pm$	0.003	0.489	$\pm$	0.003
M_V	11.18	$\pm$	0.30	11.28	$\pm$	0.30
$K(\text{km s}^{-1})$	100.54	$\pm$	0.31	101.29	$\pm$	0.31
$R(R_{\odot})$	0.453	$\pm$	0.060	0.452	$\pm$	0.050
T_{eff}^a (K)	3500			3395		

^aThe component A temperature is based on that expected of an M3V star (Cox, 2000).




Fig. 1.— Mass-Radius relation showing the M-dwarf component parameters (triangles). Our binary system is represented by squares. We also represent some other M dwarf stars, whose companions are F or G MS stars (Bouchy et al. 2005; Pont et al. 2005). We show theoretical models from Baraffe (1998), indicating an age of 10 (continuous), 5 (dash-dotted) and 1 Gyr (dashed), corresponding to $[Z]=0$, $Y=0.275$.




Fig. 2.— Phased radial velocity curve of TrES-Her0-07621 with residuals; data observed by the HET. A total of four nights were obtained, the system was observed for one hour every night at 20 minutes per exposure. The Doppler displacement was measured in two different wavelengths regions and at $H\alpha$.




Fig. 3.— Observed light curve and fitting results. The top panel shows the reconstructed lightcurve (continuous line) using the parameters obtained by fitting the observed lightcurve without the out-of-eclipse variations (small crosses, see section Results.). We have indicated phase = 0 and 1 (center of secondary eclipse) by dashed lines. The bottom panel shows the residuals (dots), with the continuous line representing the rms of residuals.